

B	I	N	G	O
u	ANTT	dress every 3-5 days	poor granulation	macerated
c	smoking/poor diet/age/diabetes	non-sterile gloves	Stage 2 - extends to epidermis & dermis	slough
Stage 3 - extends to subcutaneous tissue	gauze	FREE	l	r
length width depth	pain swelling fever reddness	elevate limb	necrosis	e
Stage 4 - extends to muscle & bone	Wound Management Plan (WMP)	non-adhesive dressing	critical colonisation	normal saline

B	I	N	G	O
venous ulceration-located lower 1/3 lower leg	pressure injury	pain swelling fever reddness	non-sterile gloves	r
Stage 4 - extends to muscle & bone	Wound Management Plan (WMP)	normal saline	necrosis	c
non-adhesive dressing	Stage 2 - extends to epidermis & dermis	FREE	gauze	ANTT
macerated	u	wound care	critical colonisation	Stage 3 - extends to subcutaneous tissue
poor granulation	debride	l	dress every 3-5 days	elevate limb

B	I	N	G	O
non-adhesive dressing	c	pain swelling fever reddness	normal saline	debride
Stage 2 - extends to epidermis & dermis	wound care	dress every 3-5 days	venous ulceration-located lower 1/3 lower leg	swab
critical colonisation	r	FREE	length width depth	poor granulation
ANTT	Wound Management Plan (WMP)	necrosis	l	exudate
u	Stage 3 - extends to subcutaneous tissue	slough	Stage 4 - extends to muscle & bone	smoking/poor diet/age/diabetes