

B	I	N	G	O
prevention	12.0 and above/4.0 and below	Severe Sepsis	given w/i 20-30 mins	antibiotics
Sepsis	pressure bag/gravity tubing	bloodstream	Activate Alert! Hosp.Policy	early recognition fast TX
more than a million	suspected /documented	Physician Order	Delay in TX	above 100.9 or below 96.8
above 20	lactate	above 90	UTI	blood culture
Sign & Symptoms of Sepsis Organ Dysfunction	SIRS	broad spectrum	vasopressors	Septic Shock

B	I	N	G	O
suspected /documented	Activate Alert! Hosp.Policy	given w/i 20-30 mins	Sepsis	bloodstream
Severe Sepsis	vasopressors	antibiotics	SIRS	early recognition fast TX
above 20	Delay in TX	Physician Order	above 100.9 or below 96.8	12.0 and above/4.0 and below
UTI	broad spectrum	prevention	Septic Shock	lactate
above 90	blood culture	more than a million	pressure bag/gravity tubing	Sign & Symptoms of Sepsis Organ Dysfunction

B	I	N	G	O
SIRS	broad spectrum	Activate Alert! Hosp.Policy	blood culture	Delay in TX
prevention	above 20	above 90	pressure bag/gravity tubing	antibiotics
more than a million	Septic Shock	Physician Order	lactate	UTI
suspected /documented	early recognition fast TX	given w/i 20-30 mins	vasopressors	Severe Sepsis
above 100.9 or below 96.8	12.0 and above/4.0 and below	bloodstream	Sepsis	Sign & Symptoms of Sepsis Organ Dysfunction