

N	U	R	S	E
Zero the transducer	300ML	Diastole	Propofol (Diprivan)	Indications for a IABP
Caregiver Needs	Sleep	Intraaortic Balloon Pumps	8-15 seconds	Phelostatic Axis
Reasons for admission to the ICU	alternate locations for pulse oximetry	FREE SPACE	Anxiety Interventions	IABP complications
ICU psychosis	Rapid Response Team	Impaired Communication	Hemodynamic Monitoring	Inications for a VAD
End of Expiration	Arterial line measurements	PA catheter complications	q 8-12 hours	Common Problems in the ICU

N	U	R	S	E
alternate locations for pulse oximetry	8-15 seconds	Propofol (Diprivan)	Caregiver Needs	Intraaortic Balloon Pumps
Diastole	q 8-12 hours	Indications for a IABP	Arterial line measurements	Phelostatic Axis
ICU psychosis	Anxiety Interventions	FREE SPACE	IABP complications	300ML
Hemodynamic Monitoring	PA catheter complications	Zero the transducer	Common Problems in the ICU	Rapid Response Team
Impaired Communication	Inications for a VAD	Reasons for admission to the ICU	Sleep	End of Expiration

N	U	R	S	E
Arterial line measurements	PA catheter complications	8-15 seconds	Indications for a VAD	Anxiety Interventions
Zero the transducer	ICU psychosis	Impaired Communication	Sleep	Indications for a IABP
Reasons for admission to the ICU	Common Problems in the ICU	FREE SPACE	Rapid Response Team	Hemodynamic Monitoring
alternate locations for pulse oximetry	Phelostatic Axis	Propofol (Diprivan)	q 8-12 hours	Diastole
IABP complications	300ML	Intraaortic Balloon Pumps	Caregiver Needs	End of Expiration