

B	I	N	G	O
The end of the bone is no longer articulated in the joint capsule	A condition in which there is increased pressure in a closed compartment	Curvature of thoracic spine	Key factor in improving patient's recovery after ortho surgery.	Helps reduce pain by reducing inflammation and swelling
Bone infection	The drugs of choice for	Inflammatory autoimmune disease of the connective tissue	A quick assessment for patient's with fractures.	Most common form of arthritis; "wear-and-tear" disease caused by injury or overuse
Treatment includes calcitonin and selective estrogen-receptor modulators	Accumulation of uric acid crystals	FREE	Exaggerated curve of lumbar spine	Five P's of compartment syndrome
Loss of bone mass	Formation of a thrombus within a deep vein	Initial pharmacotherapy goals for osteoarthritis	Dorsal flexing of the foot to assess presence of DVT	Increased risk for hypovolemic shock
Blockage in one or more arteries of the lung	Lateral deviation of spine	Care of bone disorder focus?	treating the pain and inflammation of acute gout	Compartment Syndrome

B	I	N	G	O
Treatment includes calcitonin and selective estrogen-receptor modulators	treating the pain and inflammation of acute gout	A quick assessment for patient's with fractures.	Formation of a thrombus within a deep vein	Bone infection
Key factor in improving patient's recovery after ortho surgery.	Five P's of compartment syndrome	Curvature of thoracic spine	Increased risk for hypovolemic shock	Accumulation of uric acid crystals
Exaggerated curve of lumbar spine	A condition in which there is increased pressure in a closed compartment	FREE	Dorsal flexing of the foot to assess presence of DVT	Initial pharmacotherapy goals for osteoarthritis
Compartment Syndrome	The end of the bone is no longer articulated in the joint capsule	Care of bone disorder focus?	Loss of bone mass	Blockage in one or more arteries of the lung
Helps reduce pain by reducing inflammation and swelling	Earliest manifestation of fat embolism	Most common form of arthritis; "wear-and-tear" disease caused by injury or overuse	Inflammatory autoimmune disease of the connective tissue	The drugs of choice for

B	I	N	G	O
Loss of bone mass	Dorsal flexing of the foot to assess presence of DVT	Treatment includes calcitonin and selective estrogen-receptor modulators	Compartment Syndrome	Key factor in improving patient's recovery after ortho surgery.
Lateral deviation of spine	Most common form of arthritis; "wear-and-tear" disease caused by injury or overuse	treating the pain and inflammation of acute gout	Formation of a thrombus within a deep vein	Accumulation of uric acid crystals
Initial pharmacotherapy goals for osteoarthritis	A quick assessment for patient's with fractures.	FREE	Exaggerated curve of lumbar spine	Curvature of thoracic spine
Earliest manifestation of fat embolism	Helps reduce pain by reducing inflammation and swelling	Blockage in one or more arteries of the lung	A condition in which there is increased pressure in a closed compartment	The end of the bone is no longer articulated in the joint capsule
The drugs of choice for	Inflammatory autoimmune disease of the connective tissue	Five P's of compartment syndrome	Increased risk for hypovolemic shock	Care of bone disorder focus?