

|                                                        |                           |                          |                                                   |                                     |
|--------------------------------------------------------|---------------------------|--------------------------|---------------------------------------------------|-------------------------------------|
| <b>B</b>                                               | <b>I</b>                  | <b>N</b>                 | <b>G</b>                                          | <b>O</b>                            |
| The most effective way to stop the spread of infection | 2 patient identifiers     | Code 1                   | Pt should have this device in reach at all times. | Frequency of turning/ repositioning |
| AC & HS                                                | Surgical V/S              | Isolation for TB patient | 3333                                              | Normal HR                           |
| S in SAGE                                              | Weigh or on _____ & _____ | NAs ROCK                 | 5 P's                                             | Coughing when eating is a sign of   |
| Foley care is done q shift to prevent                  | Routine V/S               | Frequency of rounding    | Code Red                                          | G in SAGE                           |
| Code 4                                                 | A in SAGE                 | NPO                      | sign of hypoglycemia/low blood sugar              | Care Channel                        |

|                                                |                                            |                                                                    |                                            |                                |
|------------------------------------------------|--------------------------------------------|--------------------------------------------------------------------|--------------------------------------------|--------------------------------|
| <b>B</b>                                       | <b>I</b>                                   | <b>N</b>                                                           | <b>G</b>                                   | <b>O</b>                       |
| Weigh or<br>on<br>_____<br>& _____             | <b>3333</b>                                | Pt should<br>have this<br>device in<br>reach at all<br>times.      | <b>AC &amp;<br/>HS</b>                     | Isolation<br>for TB<br>patient |
| <b>Code 1</b>                                  | sign of<br>hypoglycemia/low<br>blood sugar | Frequency<br>of turning/<br>repositioning                          | <b>A in<br/>SAGE</b>                       | Normal<br>HR                   |
| Foley care<br>is done q<br>shift to<br>prevent | <b>5 P's</b>                               | <b>NAs<br/>ROCK</b>                                                | Coughing<br>when<br>eating is a<br>sign of | 2 patient<br>identifiers       |
| <b>Code<br/>Red</b>                            | <b>NPO</b>                                 | The most<br>effective<br>way to stop<br>the spread<br>of infection | <b>Care<br/>Channel</b>                    | Routine<br>V/S                 |
| Frequency<br>of<br>rounding                    | <b>G in<br/>SAGE</b>                       | <b>S in<br/>SAGE</b>                                               | <b>Surgical<br/>V/S</b>                    | <b>Code 4</b>                  |

|                                                                    |                                                |                                                               |                                            |                                           |
|--------------------------------------------------------------------|------------------------------------------------|---------------------------------------------------------------|--------------------------------------------|-------------------------------------------|
| <b>B</b>                                                           | <b>I</b>                                       | <b>N</b>                                                      | <b>G</b>                                   | <b>O</b>                                  |
| A in<br>SAGE                                                       | NPO                                            | 3333                                                          | G in<br>SAGE                               | 5 P's                                     |
| The most<br>effective<br>way to stop<br>the spread<br>of infection | Foley care<br>is done q<br>shift to<br>prevent | Frequency<br>of<br>rounding                                   | Surgical<br>V/S                            | Frequency<br>of turning/<br>repositioning |
| S in<br>SAGE                                                       | Care<br>Channel                                | NAs<br>ROCK                                                   | Routine<br>V/S                             | Code<br>Red                               |
| Weigh or<br>on<br><br>_____ & _____                                | Normal<br>HR                                   | Pt should<br>have this<br>device in<br>reach at all<br>times. | sign of<br>hypoglycemia/low<br>blood sugar | Code 1                                    |
| Coughing<br>when<br>eating is a<br>sign of                         | 2 patient<br>identifiers                       | Isolation<br>for TB<br>patient                                | AC &<br>HS                                 | Code 4                                    |