

SEPSIS

Pressure injury prevention	Ask for help from peers to stay with the patient	Delivery oxygen when Spo2 drops	MAP should be above 65mmHg	Inform Doctor when blood results are back
Call doctor to come immediately without describing situation because is urgent	Pathway and other documentations	Insert catheter when needed	Ask patient to calm down and explain that nothing is happening	Not inform family until the symptoms are well managed
Roughly observations not required because ABX already given	Blood sugar check is not important in this case	Fluid balance monitor: output should be monitored closely	Given antibiotics as charted	Mechanical ventilation required to reduce work of breathing
Go to the next patient and ignore patients complain	Lab investigations as Doctors orders	Encourage more fluid intake when no urine output for 3-4 hours	Inform family and care with adequate education and updates	Blood sugar check required
MAP should be above 80mmHg	Call doctor using ISBAR handover	Sent blood culture	Call for blood test with Doctors order	Fluid balance monitor: Input should be monitored closely

SEPSIS

Roughly observations not required because ABX already given	Blood sugar check is not important in this case	Not inform family until the symptoms well managed	Ask patient to calm down and explain that nothing is happening	Blood sugar check required
Call for blood test with Doctors order	Mechanical ventilation required to reduce work of breathing	Fluid balance monitor: output should be monitored closely	Inform Doctor when blood results are back	Lab investigations as Doctors orders
Call doctor to come immediately without describing situation because is urgent	Delivery oxygen when Spo2 drops	Insert catheter when needed	Inform family and career with adequate education and updates	Pressure injury prevention
Go to the next patient and ignore patients complain	Call doctor using ISBAR handover	Pathway and other documentations	Encourage more fluid intake when no urine output for 3-4 hours	MAP should above 65mmHg
MAP should above 80mmHg	Sent blood culture	Fluid balance monitor: Input should be monitor closely	Ask for help from peers to stay with the patient	Given antibiotics as charted

SEPSIS

Mechanical ventilation required to reduce work of breathing	Inform family and care with adequate education and updates	Call doctor using ISBAR handover	Call for blood test with Doctors order	Encourage more fluid intake when no urine output for 3-4 hours
Blood sugar check required	Fluid balance monitor: Input should be monitor closely	Call doctor to come immediately without describing situation because is urgent	Fluid balance monitor: output should be monitored closely	Pressure injury prevention
Go to the next patient and ignore patients complain	MAP should above 80mmHg	Insert catheter when needed	Blood sugar check is not important in this case	Pathway and other documentations
Roughly observations not required because ABX already given	Lab investigations as Doctors orders	Sent blood culture	Ask patient to calm down and explain that nothing is happening	Ask for help from peers to stay with the patient
Given antibiotics as charted	Not inform family until the symptoms well managed	MAP should above 65mmHg	Delivery oxygen when Spo2 drops	Inform Doctor when blood results are back