

U	I	C	e	r
swab	ANTT	dress every 3-5 days	pain swelling fever reddness	necrosis
Wound Management Plan (WMP)	smoking/poor diet/age/diabetes	non-sterile gloves	Stage 2 - extends to epidermis & dermis	slough
venous ulceration-located lower 1/3 lower leg	Stage 3 - extends to subcutaneous tissue	gauze	length width depth	pressure injury
macerated	normal saline	elevate limb	non-adhesive dressing	Stage 4 - extends to muscle & bone
poor granulation	critical colonisation	wound care	debride	exudate

U	I	C	e	r
venous ulceration-located lower 1/3 lower leg	Stage 3 - extends to subcutaneous tissue	slough	Stage 2 - extends to epidermis & dermis	Stage 4 - extends to muscle & bone
debride	pressure injury	gauze	necrosis	normal saline
Wound Management Plan (WMP)	dress every 3-5 days	non-sterile gloves	non-adhesive dressing	swab
macerated	critical colonisation	smoking/poor diet/age/diabetes	elevate limb	pain swelling fever reddness
poor granulation	wound care	exudate	ANTT	length width depth

U	I	C	e	r
pressure injury	non-adhesive dressing	critical colonisation	debride	elevate limb
Stage 4 - extends to muscle & bone	exudate	Wound Management Plan (WMP)	gauze	swab
macerated	poor granulation	non-sterile gloves	Stage 3 - extends to subcutaneous tissue	smoking/poor diet/age/diabetes
venous ulceration-located lower 1/3 lower leg	normal saline	wound care	Stage 2 - extends to epidermis & dermis	ANTT
length width depth	slough	pain swelling fever reddness	dress every 3-5 days	necrosis