

U	I	C	e	r
u	ANTT	dress every 3-5 days	poor granulation	macerated
C	smoking/poor diet/age/diabetes	non-sterile gloves	Stage 2 - extends to epidermis & dermis	slough
Stage 3 - extends to subcutaneous tissue	gauze	I	r	length width depth
pain swelling fever reddness	elevate limb	necrosis	e	Stage 4 - extends to muscle & bone
Wound Management Plan (WMP)	non-adhesive dressing	critical colonisation	normal saline	venous ulceration-located lower 1/3 lower leg

U	I	C	e	r
pressure injury	pain swelling fever reddness	non-sterile gloves	r	Stage 4 - extends to muscle & bone
Wound Management Plan (WMP)	normal saline	necrosis	c	non-adhesive dressing
Stage 2 - extends to epidermis & dermis	gauze	ANTT	macerated	u
wound care	venous ulceration-located lower 1/3 lower leg	critical colonisation	Stage 3 - extends to subcutaneous tissue	poor granulation
debride	I	dress every 3-5 days	elevate limb	swab

U	I	C	e	r
critical colonisation	r	length width depth	poor granulation	ANTT
venous ulceration-located lower 1/3 lower leg	dress every 3-5 days	Wound Management Plan (WMP)	necrosis	I
exudate	u	Stage 3 - extends to subcutaneous tissue	slough	Stage 2 - extends to epidermis & dermis
Stage 4 - extends to muscle & bone	debride	smoking/poor diet/age/diabetes	gauze	macerated
pain swelling fever reddness	c	e	normal saline	non-adhesive dressing