

B	I	N	G	O
What type of diet is recommended for pt with a K level of 6.0 mEq	A ruptured achilles tendon is an adverse effect of which medication	Lab result that indicates possible prostate cancer	What is a priority nursing intervention for a pt scheduled for an Intravenous Pyelogram	Procedure that requires post op monitoring for possible bleeding on the bed linens
Hypokalemia	What organism is the most common cause of a UTI	Subjective Symptom of ESRD	Chronic anemia is often diagnoses in patient with which renal disease	Urinary diversion
What nursing intervention is necessary to determine type of calculi a pt has	Which hormone/medication is prescribed for a pt with prostate cancer	FREE SPACE	Complication of PD	Nitrofurantoin medication is prescribe for which urinary system diagnosis
Pain in the flank area is a sx of which renal complication	What is a sign of UTI in a geriatric patient	Permanent HD Access	Lab test that will determine type of antiinfective that will be administed for an infection	Procedure that the urologist will order to look for tumor calculi or inflammation of the bladder
Lab result that will help determine the correct dose of medication for a patient with diagnosis of renal failure	Hypernatremia	Uncontrolled DM can lead to which progressive and irreversible disease	which elevated lab values are monitored for a pt with decreased renal function	Hypomagnesium

B	I	N	G	O
Which hormone/medication is prescribed for a pt with prostate cancer	Chronic anemia is often diagnoses in patient with which renal disease	What is a priority nursing intervention for a pt scheduled for an Intravenous Pyelogram	Hypokalemia	Subjective Symptom of ESRD
Lab result that indicates possible prostate cancer	which elevated lab values are monitored for a pt with decreased renal function	Procedure that requires post op monitoring for possible bleeding on the bed linens	Hypernatremia	Urinary diversion
Pain in the flank area is a sx of which renal complication	Complication of PD	FREE SPACE	Nitrofurantoin medication is prescribe for which urinary system diagnosis	A ruptured achilles tendon is an adverse effect of which medication
Lab test that will determine type of antiinfective that will be administed for an infection	Uncontrolled DM can lead to which progressive and irreversible disease	What type of diet is recommended for pt with a K level of 6.0 mEq	Hypomagnesium	What is a sign of UTI in a geriatric patient
Permanent HD Access	Procedure that the urologist will order to look for tumor calculi or inflammation of the bladder	What nursing intervention is necessary to determine type of calculi a pt has	What organism is the most common cause of a UTI	Lab result that will help determine the correct dose of medication for a patient with diagnosis of renal failure

B	I	N	G	O
Hypernatremia	Uncontrolled DM can lead to which progressive and irreversible disease	Chronic anemia is often diagnoses in patient with which renal disease	Procedure that the urologist will order to look for tumor calculi or inflammation of the bladder	Complication of PD
What type of diet is recommended for pt with a K level of 6.0 mEq	Pain in the flank area is a sx of which renal complication	Permanent HD Access	What organism is the most common cause of a UTI	Procedure that requires post op monitoring for possible bleeding on the bed linens
What nursing intervention is necessary to determine type of calculi a pt has	Hypomagnesium	FREE SPACE	What is a sign of UTI in a geriatric patient	Lab test that will determine type of antiinfective that will be administed for an infection
Which hormone/medication is prescribed for a pt with prostate cancer	Urinary diversion	What is a priority nursing intervention for a pt scheduled for an Intravenous Pyelogram	which elevated lab values are monitored for a pt with decreased renal function	Lab result that indicates possible prostate cancer
Nitrofurantoin medication is prescribe for which urinary system diagnosis	A ruptured achilles tendon is an adverse effect of which medication	Subjective Symptom of ESRD	Hypokalemia	Lab result that will help determine the correct dose of medication for a patient with diagnosis of renal failure